

# Pair-distribution function analysis of the structural valence transition in $\text{Cp}_2^*\text{Yb}(4,4'\text{-Me}_2\text{-bipy})$

C H Booth<sup>1</sup>, E D Bauer<sup>2</sup>, E D Bözin<sup>3</sup>, S J L Billinge<sup>3</sup>, M D Walter<sup>1,4</sup>  
and R A Andersen<sup>1,4</sup>

<sup>1</sup>Chemical Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

<sup>2</sup>Materials Physics and Applications Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA

<sup>3</sup>Department of Physics, Columbia University, New York, New York 10027, USA

<sup>4</sup>Department of Chemistry, University of California, Berkeley, California 94720, USA

E-mail: [chbooth@lbl.gov](mailto:chbooth@lbl.gov)

**Abstract.** The  $\text{Cp}_2^*\text{Yb}(L)$  class of compounds, where  $\text{Cp}$ =pentamethylcyclopentadienyl =  $\text{C}_5\text{Me}_5$  and  $L$  is either a tertbutyldiazadiene or bipy = 2,2'-bipyridyl related ligand, have provided excellent analogies to the Kondo state on the nanoscale.  $\text{Cp}_2^*\text{Yb}(4,4'\text{-Me}_2\text{-bipy})$  furthers this analogy by demonstrating a valence transition as the sample is cooled below 200 K. Here, pair-distribution function (PDF) analysis of x-ray powder diffraction data demonstrate that the  $\text{Cp}_2^*\text{Yb}(4,4'\text{-Me}_2\text{-bipy})$  molecule is virtually unchanged through the valence transition. However, the molecule's stacking arrangement is altered through the valence transition.


In 1989, Neumann and Fulde [1] conjectured that there could be a molecular analogue to the Kondo effect in certain Ce- and Yb-based organometallic molecules. The theory of such effects (see Ref. [2] and references therein) where the traditional role of many-body interactions in Kondo systems are reduced to, essentially, including specific configuration interactions (CIs) to the single-electron theory using, for instance, the Complete Active Space Self-Consistent Field (CASSCF) approach. According to such methods, intermediate valence (IV) occurs in the ground state when open and closed shell singlet configurations mix, thereby lowering the combined open-shell singlet state below the triplet state [2]. For instance, in  $\text{Cp}_2^*\text{Yb}(\text{bipy})$ , where  $\text{Cp}^*$  = pentamethylcyclopentadienyl =  $\text{C}_5\text{Me}_5$ , Me = methyl =  $\text{CH}_3$ , and bipy = 2,2'-bipyridine =  $(\text{C}_5\text{H}_4\text{N})_2$ , Density Functional Theory (DFT) calculations indicate a triplet  $f_{\uparrow}^{13}\pi_{\uparrow}^{*1}$  in the ground state [2]. However, the higher-order CASSCF calculations indicate that the open-shell  $f_{\uparrow}^{13}\pi_{\downarrow}^{*1}$  configuration interacts with, primarily, the closed-shell  $f_{\uparrow,\downarrow}^{14}\pi^{*0}$  configuration, to form an open-shell singlet below the triplet configuration, with an  $f$ -hole occupancy of about  $n_f = 0.8$ , consistent with Yb  $L_{\text{III}}$ -edge x-ray absorption near-edge structure (XANES) experiments [2, 3]. In analogy to the Kondo effect, the singlet interactions are made possible both by symmetry and due to the more extended nature of the  $f$ -orbital in Yb and the  $\pi^*$  orbital in the aromatic bipy ligand, which plays the role of a quantum-confined metallic band in an intermetallic nanoparticle [4, 5]. The Kondo analogy has worked very well for guiding many other aspects of the research [6, 2]. Such results therefore have broad implications for fields such as solid-state physics, materials science, and organometallic chemistry, and may have future implications for nanoscale devices, which have spurred recent interest in understanding the Kondo effect on the nanoscale.

The Me-substituted  $\text{Cp}_2^*\text{Yb}(\text{bipy})$  complexes demonstrate one of the more dramatic differences between bulk and nanoscale Kondo behavior in that the first excited state need not be a triplet. However, the  $\text{Cp}_2^*\text{Yb}(4,4'\text{-Me}_2\text{-bipy})$  molecule also undergoes a first-order +10% valence transition when cooling below 200 K [7, 3]. Here, we address the low temperature structure of the molecule using the pair-distribution function (PDF) technique to determine whether the molecule is isomorphic through the transition, in analogy to the isomorphic valence transition in  $\text{YbInCu}_4$  [8] and the  $\alpha$ - $\gamma$  transition in elemental cerium [9, 10].


Synthesis details for the  $\text{Cp}_2^*\text{Yb}(4,4'\text{-Me}_2\text{-bipy})$  complex are reported in Ref. [7]. In preparation for the x-ray scattering experiments, the air-sensitive single crystals were powdered in an inert atmosphere glove box. The finely powdered sample was packed into a 5 mm diameter hole in a 1 mm thick aluminum plate, and then sealed between aluminized mylar windows. The sample was then shipped to the Advanced Photon Source (APS) at Argonne National Laboratory in a nitrogen-filled jar. The sample was removed from the jar and then placed into a displac just prior to measurement at the 6ID-D beamline of MuCAT. Measurements were made at many temperatures between 50 K and 300 K. X-ray powder-diffraction data were collected using the rapid acquisition pair distribution function method [11]. The experiments were conducted using synchrotron x-rays with a wavelength of 0.126340 Å using a circular image plate camera (Mar345) 345 mm in diameter. The camera was mounted orthogonally to the beam path with a sample-to-detector distance of 240.4640 mm. Other details of data collection and analysis are as described previously [12]. The corrected total scattering structure function,  $S(Q)$ , was obtained using standard corrections [13] with the program PDFGETX2 [14]. Finally, the PDF was obtained by Fourier transformation of  $S(Q)$  according to  $G(r) = \frac{2}{\pi} \int_0^{Q_{\text{max}}} Q[S(Q) - 1] \sin(Qr) dQ$ , where  $Q$  is the magnitude of the scattering vector. A  $Q_{\text{max}} = 15.0 \text{ \AA}^{-1}$  was used. Fig. 2 shows  $G(r)$  for several selected temperatures. Fits are performed using the PDFGUI analysis package [15].

From room temperature to just above the transition, the changes in the PDF data from temperature to temperature are small, consistent with thermally-induced variations. At the transition, a clearly resolved change occurs, as illustrated in Fig. 2, where in the 10 K above the transition the data overlap well (Fig. 2a), yet in the next 10 K (Fig. 2b), the changes in the local structure are easily discernible, especially starting at about 9 Å. However, below 9 Å, the local structure is nearly unaffected. The intermolecular Yb-Yb distance is about 9 Å in these crystals. We can therefore conclude simply from these data and with no fitting that the main structural change occurs in the molecular stacking.


There remain two unanswered questions: (1) can we resolve an actual change in the molecule's morphology, and (2) what is the change in the packing arrangement? To answer these questions, a detailed fit should be performed. Unfortunately, attempts to fit such models to the low temperature data, even including a wide variety of constraints, did not produce satisfactory



**Figure 1.** The  $\text{Cp}_2^*\text{Yb}(4,4'\text{-Me}_2\text{-bipy})$  molecule and the position numbering scheme for bipyridine. The  $\text{Cp}_2^*\text{Yb}(4,4'\text{-Me}_2\text{-bipy})$  molecule crystallizes into the  $P2_1/c$  space group above 200 K.



**Figure 2.** The distribution function at two temperatures 10 K apart (a) above and (b) below the transition.



**Figure 3.** (a) Data and fit to the single crystal structure. (b) Fit from (a) together with the same fit with the 4,4'-Me<sub>2</sub> removed. (c) Fit from (a) together with a calculation of Cp<sub>2</sub>\*Yb(5,5'-Me<sub>2</sub>-bipy), in the *Pbca* space group, using the same thermal parameters.

results. It is possible to fit the high-temperature-phase data to the nominal crystal structure [7] by only allowing a single mean-squared displacement parameter  $u^2$  for each atomic species, the three lattice parameters, a scale factor, and a damping factor to vary. Such a fit is shown in Fig. 3a. Clearly, the high-temperature-phase data represents the structure from the much more accurate single-crystal diffraction results well.

Fortunately, we can demonstrate the sensitivity of the data (if not the fit) to possible changes in the molecule's morphology through the transition using these fit results. A relatively small change in the morphology, at least as far as the x-ray scattering is concerned, would be to eliminate a C atom from the molecule. In Fig. 3b, the effect of removing the Me carbons at the 4,4' sites on the bipy is illustrated by taking the fit result shown in Fig. 3a and simply deleting these Me groups. The  $G(r)$  is nearly unaffected, except at the Yb-Me distance of about 7 Å. This example, of course, illustrates the change from two carbon atoms; in any case, there is no change observed on this scale below 9 Å in Fig. 2b. Consequently, we conclude that the molecule's morphology is unaffected by the valence transition. Changes in the bond lengths due to the change in valence can be observed in these data, but are more accurately presented in fits to extended x-ray absorption fine-structure (EXAFS) data described elsewhere [3].

Although the molecule has the same morphology in the high and low temperature phases, it is possible, if not likely, that the change in packing produces a crystal structure with a different symmetry than  $P2_1/c$ . A likely alternative is a *Pbca* structure, such as occurs for Cp<sub>2</sub>\*Yb(bipy), Cp<sub>2</sub>\*Yb(5,5'-Me<sub>2</sub>-bipy), and others [3]. As mentioned above, fits using the *Pbca* structure are inconclusive, as illustrated by the calculation in Fig. 3c, where the structure of Cp<sub>2</sub>\*Yb(5,5'-Me<sub>2</sub>-bipy) is used with thermal constants generated from the fit shown in Fig. 3a to Cp<sub>2</sub>\*Yb(4,4'-Me<sub>2</sub>-bipy) data. Clearly, the agreement is very poor. Attempts to fit these data

using Rietveld refinement also failed to produce a unique answer, since the data can be fit using either space group if enough parameters are allowed to vary.

The molecule itself is therefore isomorphous through the valence transition, but based on the changes in packing, the most likely conclusion regarding the crystal structure is that it is not isomorphous through the transition. It remains unclear whether the valence transition is driven by the Kondo effect or not. As the low-temperature structure may well be in the *Pbca* space group, the change in structure could be driven by steric effects. The observed valence transition would then be an indication that intermolecular, van der Waals interactions are important in these molecules, although we note that CASSCF calculations on isolated molecules obtain an  $n_f$  that is generally well within 10% of the measured values. The changing intermolecular interaction could, in any case, have an effect on the overall  $\pi^*$  charge density, and so the change in valence could be a manifestation of a charge-density induced change to the  $f$ -orbital coupling. This scenario would be very similar to one suggested for YbInCu<sub>4</sub>, except that in that case the change in density is thought to be due to a sharp edge to a band near the Fermi level [16].

Another possibility is that intermolecular effects are insignificant, and the valence transition is somehow analogous to that which occurs in the  $\alpha$ - $\gamma$  transition in cerium [9, 10]. Of course, even here, there remains a controversy regarding the basic physics underlying the volume collapse in elemental cerium. The two general possibilities at present are the so-called Kondo Volume Collapse model (KVM) [17], and the Mott transition model [18]. With regard to the role of the Kondo effect, the KVM achieves a volume change by exploiting nonlinearities in the pressure/volume relationship in the Kondo impurity model [17]. Although the possibility of a Mott transition exists in cerium, it is unlikely that there is some local equivalent possibility in Cp<sub>2</sub>\*Yb(4,4'-Me<sub>2</sub>-bipy) given the observed IV in both the high temperature and the low temperature phases. Therefore, CI-induced IV behavior appears to play an important role in generating the valence transition in Cp<sub>2</sub>\*Yb(4,4'-Me<sub>2</sub>-bipy), either through structural-induced changes in charge density or through a KVM-like mechanism.

## Acknowledgments

Work at Lawrence Berkeley National Laboratory was supported by the Director, Office of Science (OS), Office of Basic Energy Sciences (OBES), of the U.S. Department of Energy (DOE) under Contract No. DE-AC02-05CH11231. Work in the Billinge group was supported in part by the National Science Foundation under Grants No. DMR-0304391 and No. DMR-0703940. Work at Los Alamos was supported under the auspices of the DOE. Use of the APS was supported by the U. S. DOE, OS, OBES, under Contract No. DE-AC02-06CH11357.

## References

- [1] Neumann C S and Fulde P 1989 *Z. Phys. B* **74** 277
- [2] Booth C H, Walter M D, Kazhdan D, Hu Y J, Lukens W W, Bauer E D, Maron L, Eisenstein O and Andersen R A 2009 *J. Am. Chem. Soc.* **131** 6480
- [3] Booth C H *et al.* Intermediate-valence tautomerism in decamethylterbocene complexes of methyl-substituted bipyridines (*Preprint* [http://lise.lbl.gov/chbooth/papers/Me\\_bipy.pdf](http://lise.lbl.gov/chbooth/papers/Me_bipy.pdf))
- [4] Thimm W B, Kroha J and von Delft J 1999 *Phys. Rev. Lett.* **82** 2143
- [5] Schlottmann P 2002 *Phys. Rev. B* **65** 024420
- [6] Booth C H, Walter M D, Daniel M, Lukens W W and Andersen R A 2005 *Phys. Rev. Lett.* **95** 267202
- [7] Walter M D, Berg D J and Andersen R A 2006 *Organometallics* **25** 3228
- [8] Felner I and Nowik I 1986 *Phys. Rev. B* **33** 617
- [9] Lawrence J M, Riseborough R S and Parks R D 1981 *Rep. Prog. Phys.* **44** 1
- [10] McMahan A K, Huscroft C, Scalettar R T and Pollock E L 1998 *J. Comput.-Aid. Mater. Des.* **5** 131
- [11] Chupas P J, Qiu X, Hanson J C, Lee P L, Grey C P and Billinge S J L 2003 *J. Appl. Crystallogr.* **36** 1342
- [12] Lin H, Božin E S, Billinge S J L, Androulakis J, Malliakas C D, Lin C H and Kanatzidis M G 2009 *Phys. Rev. B* **80** 045204

- [13] Egami T and S J L Billinge 2003 *Underneath the Bragg peaks: Structural analysis of complex materials* (Amsterdam: Pergamon)
- [14] Qiu X, Thompson J W and Billinge S J L 2004 *J. Appl. Crystallogr.* **37** 678
- [15] Farrow C L, Juhás P, Liu J W, Bryndin D, Božin E S, Bloch J, Proffen T and Billinge S J L 2007 *J. Phys.: Condens. Matter* **19** 335219
- [16] Figueroa E, Lawrence J M, Sarrao J L, Fisk Z, Hundley M F and Thompson J D 1998 *Solid State Commun.* **106** 347
- [17] Allen J W and Martin R M 1982 *Phys. Rev. Lett.* **49** 1106
- [18] Johansson B 1974 *Phil. Mag.* **30** 469